

ZIG-ZAG CHAINS AND METRIC EQUIVALENCES BETWEEN ULTRAMETRIC SPACES

ÁLVARO MARTÍNEZ-PÉREZ

ABSTRACT. We study the classification of ultrametric spaces based on their small scale geometry (uniform homeomorphism), large scale geometry (coarse equivalence) and both (all scale uniform equivalences). We prove that these equivalences can be characterized with parallel constructions using a combinatoric tool called *common zig-zag chain*.

CONTENTS

1. Introduction	1
2. Expansion functions	4
3. All scale uniform equivalences	5
4. Coarse equivalences	9
5. Uniform homeomorphisms	11
6. Towers and admissible morphisms	14
References	16

Keywords: Ultrametric, chain, end space, coarse, uniform homeomorphism, zig-zag chain.

MSC: primary 18B30, 37F20; secondary 54E35.

1. INTRODUCTION

When one defines continuity for a function on a metric space, one neglects a great deal of the information contained in the metric focusing on the small scale structure. In fact, if d is a metric then so it is $d' = \min\{d, 1\}$ and this change won't affect continuity nor the topology of the space.

The dual situation appears with bornologous maps, where we pay attention only to the large scale geometry. If we consider the metric $d' = \max\{d, 1\}$ all the topology of the space is lost, but we still keep all its large scale properties. For a further development of this, see [9].

Thus, uniform category and coarse category are partial and somehow dual aspects of the whole picture. To depict both scales we use *all scale uniform maps*, which are both, uniformly continuous and bornologous.

In this paper, we consider three categories of ultrametric spaces.

- $\mathcal{C}_1$: Complete ultrametric spaces and all scale uniform maps.
- $\mathcal{C}_2$: Ultrametric spaces and surjective bornologous multi-maps.
- $\mathcal{C}_3$: Complete ultrametric spaces and uniformly continuous maps.

Partially supported by MTM 2006-00825.

The idea is to characterize equivalences in these three categories using the same combinatorial technique and the same arguments, presenting categories $\mathcal{C}_2$ and $\mathcal{C}_3$ as partial representations of the geometry in $\mathcal{C}_1$.

Let us recall here the definition and basic properties of ultrametric spaces.

Definition 1.1. *If (X, d) is a metric space and $d(x, y) \leq \max\{d(x, z), d(z, y)\}$ for all $x, y, z \in X$, then d is an ultrametric and (X, d) is an ultrametric space.*

- Lemma 1.2.** (a) *Any point of a ball is a center of the ball.*
 (b) *If two balls have a common point, one is contained in the other.*
 (c) *The diameter of a ball is less than or equal to its radius.*
 (d) *In an ultrametric space, all triangles are isosceles with at most one short side.*
 (e) $S_r(a) = \bigcup_{x \in S_r(a)} B_{<r}(x)$.
 (f) *The spheres $S_r(a)$ ($r > 0$) are both open and closed.* ■

There is a well known correspondence between ultrametric spaces and trees. In an ultrametric space, for any pair of intersecting balls one will contain the other and hence, considering partitions of the space with shrinking diameter we obtain a branching process which can be modeled by a tree.

In the bounded case, B. Hughes establishes some categorical equivalences in [4], which capture the geometry of trees at infinity and local geometry of ultrametric spaces.

From a more topological point of view, in [7], it is proved that there is categorical equivalence between complete ultrametric spaces of diameter ≤ 1 with uniformly continuous maps and rooted geodesically complete $\mathbb{R}$ -trees with classes of rooted, continuous and metrically proper maps. Hence, uniform homeomorphisms between bounded ultrametric spaces, can be characterized by some kind of metrically proper homotopy equivalence between the trees. The technique to do this uses a function called *modulus of continuity* which is associated to any uniformly continuous map. This idea is used here in a generalized way defining what will be called *expansion function*.

In [1] Taras Banach and Ihor Zarichnyy characterize coarse equivalences of homogeneous ultrametric spaces by some intrinsic invariant of the spaces called sharp entropy. They do this using induction on partially ordered sets called towers. In a slightly different approach, these objects are treated here as chains instead of as ordered sets.

Trees are also related to chains and inverse sequences. In [8] it is proved an equivalence of categories between inverse sequences and rooted geodesically complete $\mathbb{R}$ -trees oriented to a geometric description of the shape category in Mardešić-Segal approach (see [6]). In this paper, we defined a functor from maps between trees to morphisms of inverse sequences related to the construction used here.

A *directed chain* (X_k, Φ_k) is a collection of sets X_k $k \in \mathbb{Z}$ and maps $\Phi_k : X_k \rightarrow X_{k+1}$ $k \in \mathbb{Z}$. The *direct limit*, $\varinjlim X_k$, is the disjoint union of the X_k 's modulo some equivalence relation $\sim$: for any pair of points $x_i \in X_i$, $x_j \in X_j$,

$$x_i \sim x_j \text{ if there is some } k > i, j \text{ such that } f_{k-1} \circ \cdots \circ f_i(x_i) = f_{k-1} \circ \cdots \circ f_j(x_j).$$

Definition 1.3. *A D -chain (X_k, Φ_k) is a collection of sets X_k $k \in \mathbb{Z}$ and surjective maps $\Phi_k : X_k \rightarrow X_{k+1}$, $k \in \mathbb{Z}$, such that $\varinjlim X_k$ is trivial.*

Hence, using all $k \in \mathbb{Z}$ we characterize the all scale uniform type of U , and if we want to focus only on the large scale or the small scale structure, we only need to restrict ourselves, roughly speaking, to one side of the chain.

Definition 1.4. A D_+ -chain (X_n, Φ_n) is a sequence of sets X_n and surjective maps $\Phi_n : X_n \rightarrow X_{n+1}$ such that $\lim_{\rightarrow} X_n$ is trivial.

Definition 1.5. A D_- -chain (X_n, p_n) is a sequence of sets X_n and surjective maps $\Phi_n : X_{n+1} \rightarrow X_n$.

Definition 1.6. Given a D -chain (X_k, Φ_k) and an increasing function $\alpha : \mathbb{Z} \rightarrow \mathbb{Z}$, the D -chain defined by the sets $X_{\alpha(k)}$ and the maps $\tilde{\phi}_k = \phi_{\alpha(k+1)-1} \circ \cdots \circ \Phi_{\alpha(k)}$ will be called an α -sub- D -chain.

Definition 1.7. Given a D_+ -chain (X_n, Φ_n) and an increasing function $\alpha : \mathbb{N} \rightarrow \mathbb{N}$, the D_+ -chain defined by the sets $X_{\alpha(n)}$ and the maps $\tilde{\phi}_n = \phi_{\alpha(n+1)-1} \circ \cdots \circ \Phi_{\alpha(n)}$ will be called an α -sub- D_+ -chain.

Definition 1.8. Given a D_- -chain (X_n, Φ_n) and an increasing function $\alpha : \mathbb{N} \rightarrow \mathbb{N}$, the D_- -chain defined by the sets $X_{\alpha(n)}$ and the maps $\tilde{\phi}_n = \phi_{\alpha(n+1)-1} \circ \cdots \circ \Phi_{\alpha(n)}$ will be called an α -sub- D_- -chain.

Remark 1.9. When there is no need to specify the map α and it is clear from the context whether we are considering D -chains, D_+ -chains or D_- -chains, we will call these just *subchains*.

The following definitions are adapted from [5].

Definition 1.10. A D -chain $(Z_k, \mathcal{V}_k)$ is a common zig-zag D -chain of the D -chains $(X_n, \Phi_n), (Y_n, \Psi_n)$ if there are increasing maps $\alpha, \beta : \mathbb{Z} \rightarrow \mathbb{Z}$ and subchains $(X_{\alpha(k)}, \tilde{\Phi}_k), (Y_{\beta(k)}, \tilde{\Psi}_k)$ with

$$Z_i = \begin{cases} X_{\alpha(\frac{i+1}{2})} & \text{if } i \text{ is odd,} \\ Y_{\beta(\frac{i}{2})} & \text{if } i \text{ is even.} \end{cases}$$

such that the following diagram commutes

$$\begin{array}{ccccccc} \longrightarrow & X_{\alpha(k-1)} & \longrightarrow & X_{\alpha(k)} & \longrightarrow & X_{\alpha(k+1)} & \longrightarrow \\ & \searrow \nu_{2k-3} & & \nearrow \nu_{2k-2} & \searrow \nu_{2k-1} & \nearrow \nu_{2k} & \searrow \nu_{2k+1} \\ \cdots & \longrightarrow & Y_{\beta(k-1)} & \longrightarrow & Y_{\beta(k)} & \longrightarrow & Y_{\beta(k+1)} \end{array}$$

Definition 1.11. A D_+ -chain $(Z_n, \mathcal{V}_n)$ is a common zig-zag D_+ -chain of the D_+ -chains $(X_n, \Phi_n), (Y_n, \Psi_n)$ if there are increasing maps $\alpha, \beta : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ and subchains $(X_{\alpha(n)}, \tilde{\Phi}_n), (Y_{\beta(n)}, \tilde{\Psi}_n)$ with

$$Z_i = \begin{cases} X_{\alpha(\frac{i+1}{2})} & \text{if } i \text{ is odd,} \\ Y_{\beta(\frac{i}{2})} & \text{if } i \text{ is even.} \end{cases}$$


such that the following diagram commutes

$$\begin{array}{ccccccc} X_{\alpha(1)} & \longrightarrow & X_{\alpha(2)} & \longrightarrow & X_{\alpha(3)} & \longrightarrow & \cdots \\ & \searrow \nu_1 & & \nearrow \nu_2 & \searrow \nu_3 & \nearrow \nu_4 & \searrow \nu_5 \\ & Y_{\beta(1)} & \longrightarrow & Y_{\beta(2)} & \longrightarrow & Y_{\beta(3)} & \longrightarrow \end{array}$$

Definition 1.12. A D_- -chain $(Z_n, \mathcal{V}_n)$ is a common zig-zag D_- -chain of the D_- -chains $(X_n, \Phi_n), (Y_n, \Psi_n)$ if there are increasing maps $\alpha, \beta: \mathbb{N} \rightarrow \mathbb{N}$ and subchains $(X_{\alpha(n)}, \tilde{\Phi}_n), (Y_{\beta(n)}, \tilde{\Psi}_n)$ with

$$Z_i = \begin{cases} X_{\alpha(\frac{i+1}{2})} & \text{if } i \text{ is odd,} \\ Y_{\beta(\frac{i}{2})} & \text{if } i \text{ is even.} \end{cases}$$

such that the following diagram commutes



The main result would be that two chains represent the same class of ultrametric space in the category $\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3$ respectively if and only if there is a common zig-zag chain (D -chain, D_+ -chain or D_- -chain) of them.

Notation 1.13. We will denote by $(X_k, \Phi_k) \sim_{z-z} (Y_k, \Psi_k)$ if there is a common zig-zag chain of the chains $(X_k, \Phi_k), (Y_k, \Psi_k)$.

This is related to some results in [1]. In their work, they consider towers as ordered sets, which is just an alternative definition for what here is called D_+ -chain. Also, we define here the end space of a chain and an ultrametric on it which is not exactly the same as they do.

Lemma 2 in [1] states:

Proposition 1.14. Let $\phi: T_1 \rightarrow T_2$ be an admissible morphism between towers T_1, T_2 . Then, the restriction $\Phi = \phi|_{[T_1]}: [T_1] \rightarrow [T_2]$ is an isomorphism. ■

Here we proof that there exists an admissible map if and only if there is a common zig-zag D_+ -chain for the D_+ -chains corresponding to the towers T_1, T_2 and that this implies a partial converse to 1.14.

2. EXPANSION FUNCTIONS

Let us recall first some definitions in coarse geometry.

A map between metric spaces $f: X \rightarrow Y$ is *metrically proper* if for any bounded set $A \in Y$, $f^{-1}(A)$ is bounded in X .

A map between metric spaces $f: X \rightarrow Y$ is *bornologous* if for every $R > 0$ there is $S > 0$ such that for any two points $x, x' \in X$ with $d(x, x') < R$, $d(f(x), f(x')) < S$.

A map is *coarse* if it is metrically proper and bornologous.

Two maps between metric spaces $f, g: X \rightarrow Y$ are *close* if $\sup_{x \in X} \{d(f(x), g(x))\} < \infty$.

A coarse map $f: X \rightarrow Y$ is a *coarse equivalence* if there is a coarse map $g: Y \rightarrow X$ such that $g \circ f$ is close to id_X and $f \circ g$ is close to id_Y . If there are such maps, then X, Y are *coarse equivalent*.

But this is not the only way to define coarse equivalence between metric spaces. In this section, in order to describe in the same terms the categories $\mathcal{C}_1, \mathcal{C}_2$ and $\mathcal{C}_3$, we will use the following definition with multi-maps, as in [1].

By a multi-map $\Phi: X \Rightarrow Y$ between two sets X, Y we understand any subset $\Phi \subset X \times Y$. For any subset $A \subset X$, by $\Phi(A) = \{y \in Y : \exists a \in A \text{ with } (a, y) \in \Phi\}$ we denote the image of A under the multi-map Φ . The inverse $\Phi^{-1}: Y \rightarrow X$ to

the multi-map Φ is the subset $\Phi^{-1} = \{(y, x) \in Y \times X : (x, y) \in \Phi\} \subset Y \times X$. For two multi-maps $\Phi : X \Rightarrow Y$, $\Psi : Y \Rightarrow Z$ the composition $\Psi \circ \Phi$ is defined as usual:

$$\Psi \circ \Phi = \{(x, z) \in X \times Z : \exists y \in Y \text{ such that } (x, y) \in \Phi \text{ and } (y, z) \in \Psi\}.$$

A multi-map is called *surjective* if $\Phi(X) = Y$ and *bijective* if $\Phi \subset X \times Y$ coincides with the graph of a bijective (single-valued) function.

Definition 2.1. *Given a multi-map $\Phi : X \Rightarrow Y$ between metric spaces, a non-decreasing function $\varrho_\Phi : J \rightarrow [0, \infty)$ with $J = [0, S]$ or $J = [0, \infty)$ is called expansion function if $\forall A \in X$ with $\text{diam}(A) \in J$, $\text{diam}(\Phi(A)) \leq \varrho_\Phi(\text{diam}(A))$.*

Definition 2.2. *A multi-map $\Phi : X \Rightarrow Y$ between metric spaces is called*

- *bornologous if there is an expansion function $\varrho_\Phi : [0, \infty) \rightarrow [0, \infty)$.*
- *an asyrmorphism if both Φ, Φ^{-1} are surjective bornologous multi-maps.*

The following characterization is contained in Proposition 2 in [1].

Proposition 2.3. *For metric spaces X, Y the following assertions are equivalent:*

- *X and Y are asyrmorphic.*
- *X and Y are coarse equivalent.* ■

Remark 2.4. *Thus, equivalences in $\mathcal{C}_2$ are, in fact, coarse equivalences of ultrametric spaces.*

Definition 2.5. Φ is called all scale uniform if there is an expansion function $\varrho_\Phi : [0, \infty) \rightarrow [0, \infty)$ such that $\varrho_\Phi(t) = 0$ and $\lim_{t \rightarrow 0} \varrho_\Phi(t) = 0$. In this case, since $\varrho_\Phi(t) = 0$, Φ is a single-valued map. If Φ^{-1} is also all scale uniform we say that X, Y are all scale uniform equivalent.

Definition 2.6. A map $f : X \rightarrow Y$ between metric spaces is uniformly continuous if $\forall \epsilon > 0$ there exists some $\delta > 0$ such that for any pair of points x, y with $d_X(x, y) < \delta$ then $d_Y(f(x), f(y)) < \epsilon$.

Proposition 2.7. A map $\Phi : X \rightarrow Y$ between metric spaces is uniformly continuous if and only if there is an expansion function $\varrho_\Phi : [0, S] \rightarrow [0, \frac{1}{2}]$ such that $\varrho_\Phi(0) = 0$ and $\lim_{t \rightarrow 0} \varrho_\Phi(t) = 0$.

Proof. The if part is obvious.

If Φ is uniformly continuous here is some $S > 0$ such that for any pair of points x, y such that $d_X(x, x') < S$ then $d_Y(f(x), f(x')) < \frac{1}{2}$. Thus, it suffices to take $\varrho_\Phi(t) := \sup_{x, x' \in X, d(x, x') \leq t} \{d(\Phi(x), \Phi(x'))\}$. □

3. ALL SCALE UNIFORM EQUIVALENCES

Given an all scale uniform map Φ and an expansion function $\varrho_\Phi : [0, \infty) \rightarrow [0, \infty)$ such that $\varrho_\Phi(t) = 0$ and $\lim_{t \rightarrow 0} \varrho_\Phi(t) = 0$, let us define $\gamma_{\varrho_\Phi} : \mathbb{Z} \rightarrow \mathbb{Z}$ as follows,

$$(1) \quad \gamma_{\varrho_\Phi}(k) := [\log_2(\varrho_\Phi(2^k))] + 1$$

where $[t]$ stands for the maximal integer less or equal than t . Hence, for all points $x, x' \in X$, if $d_X(x, x') \leq 2^k$ then $d_Y(f(x), f(x')) \leq 2^{\gamma_{\varrho_\Phi}(k)}$ and γ_{ϱ_Φ} is non-decreasing. Then $\lim_{k \rightarrow -\infty} \gamma_{\varrho_\Phi}(k) = -\infty$ since Φ is uniformly continuous and we may assume, with no loss of generality, that $\lim_{t \rightarrow \infty} \varrho_\Phi(t) = \infty$ and therefore $\lim_{n \rightarrow \infty} \gamma_{\varrho_\Phi}(n) = \infty$.

Remark 3.1. If Φ is an all scale uniform equivalence between unbounded metric spaces then necessarily $\lim_{t \rightarrow \infty} \varrho_\Phi(t) = \infty$ since Φ^{-1} is a bornologous surjective map.

There is a correspondence between complete ultrametric spaces and D -chains. Let U be an ultrametric space. For each $k \in \mathbb{Z}$ let X_k be the partition of U in balls of radius 2^k . Let $\Phi_k: X_k \rightarrow X_{k+1}$ the map canonically induced by the inclusion for any $k \in \mathbb{Z}$. (X_k, Φ_k) will be called the *extended chain associated to U* . Conversely, given a D -chain we can obtain an ultrametric space as follows.

Let us define the end space as follows:

$$\text{end}(X_k, \Phi_k) := \{(x_k)_{k \in \mathbb{Z}} \mid x_k \in X_k \text{ and } \Phi(x_k) = x_{k+1}\},$$

and let us define the metric

$$D((x_k), (y_k)) = 2^{k_0} \text{ where } k_0 = \min\{k : x_k = y_k\}.$$

D is well defined since $\lim_{\rightarrow} (X_k, \Phi_k)$ is trivial, and clearly, D is an ultrametric.

Proposition 3.2. If (U, d) is a complete ultrametric space and (X_k, Φ_k) is the D -chain associated to U , then (U, d) and $(\text{end}(X_k, \Phi_k), D)$ are bi-Lipschitz equivalent. In particular, they are all scale uniform equivalent.

Proof. First, note that if (U, d) is complete there is a bijection $i: U \rightarrow \text{end}(X_k, \Phi_k)$.

By the properties of the ultrametric, $d(x, y) \leq 2^k$ if and only if the points are in the same ball in the partition X_k . Hence, $d(x, y) \leq D(i(x), i(y)) \leq 2d(x, y)$. $\square$

Definition 3.3. A morphism of D -chains $(f_k, \sigma): (X_k, \Phi_k) \rightarrow (Y_k, \Psi_k)$ consists of a non-decreasing function $\sigma: \mathbb{Z} \rightarrow \mathbb{Z}$ such that $\lim_{k \rightarrow -\infty} \sigma(k) = -\infty$ and $\lim_{k \rightarrow \infty} \sigma(k) = \infty$, and maps $f_k: X_k \rightarrow Y_{\sigma(k)}$ such that the following diagram commutes:

$$\begin{array}{ccccccc} X_{k-1} & \longrightarrow & X_k & \longrightarrow & X_{k+1} & \longrightarrow & \\ \searrow f_{k-1} & & \searrow f_k & & \searrow f_{k+1} & & \\ \longrightarrow & Y_{\sigma(k-1)} & \longrightarrow & \cdots & \longrightarrow & Y_{\sigma(k)} & \longrightarrow \cdots \longrightarrow Y_{\sigma(k+1)} \end{array}$$

Remark 3.4. Notice that although in this definition $\sigma(k)$ and $\sigma(k+1)$ may be the same for some k , using that $\lim_{k \rightarrow -\infty} \sigma(k) = -\infty$ and $\lim_{k \rightarrow \infty} \sigma(k) = \infty$ then for some function α , we may assume that σ is increasing when restricted to the α -subchain.

Lemma 3.5. If (X_k, Φ_k) is a D -chain, $\alpha: \mathbb{Z} \rightarrow \mathbb{Z}$ is an increasing map and $(X_{\alpha(k)}, \tilde{\Phi}_k)$ is the α -subchain, then $\text{end}(X_k, \Phi_k)$ is all scale uniform equivalent to $\text{end}(X_{\alpha(k)}, \tilde{\Phi}_k)$.

Proof. Consider the canonical map $i: \text{end}(X_k, \Phi_k) \rightarrow \text{end}(X_{\alpha(k)}, \tilde{\Phi}_k)$.

Let us define the function $\Lambda: \mathbb{Z} \rightarrow \mathbb{Z}$ such that $\Lambda(z) = \min\{k : \alpha(k) \geq z\}$. Since α is increasing, it follows that Λ is non-decreasing, $\lim_{z \rightarrow -\infty} \Lambda(z) = -\infty$ and $\lim_{z \rightarrow \infty} \Lambda(z) = \infty$. Now, consider the function $\Gamma: [0, \infty) \rightarrow [0, \infty)$ such that $\Gamma(0) = 0$ and for any $x \in (2^{k-1}, 2^k]$, $\Gamma(x) = 2^{\Lambda(k)}$. Clearly Γ is non-decreasing, $\lim_{x \rightarrow 0} \Gamma(x) = 0$ and $\lim_{x \rightarrow \infty} \Gamma(x) = \infty$. For any two end points $(x_k), (y_k) \in \text{end}(X_k, \Phi_k)$ with $D((x_k), (y_k)) = 2^{k_0}$ the distance between their correspondent subsequences $i(x_k) = (x_{\alpha(k)}), i(y_k) = (y_{\alpha(k)})$ is exactly $D'((x_{\alpha(k)}), (y_{\alpha(k)})) = \Gamma(2^{k_0}) = 2^{\Lambda(k_0)}$. Then Γ is an extension function for $i: \text{end}(X_k, \Phi_k) \rightarrow \text{end}(X_{\alpha(k)}, \tilde{\Phi}_k)$ and i is an all scale uniform map. A similar argument works for i^{-1} . $\square$

Proposition 3.6. *Consider two complete ultrametric spaces U_1, U_2 and let $(X_k, \Phi_k), (Y_k, \Psi_k)$ be their associated D -chains. Then, there is an all scale uniform map $f: U_1 \rightarrow U_2$ if and only if there is a morphism of D -chains $(f_k, \sigma): (X_k, \Phi_k) \rightarrow (Y_k, \Psi_k)$.*

Proof. If there is an all scale uniform map $\Phi: U_1 \rightarrow U_2$, consider the map $\gamma_{\varrho_\Phi}: \mathbb{Z} \rightarrow \mathbb{Z}$ from (1). Let us consider $\sigma = \gamma_{\varrho_\Phi}$. Then, Φ induces maps $f_k: X_k \rightarrow Y_{\sigma(k)}$ canonically as follows: by construction, any point $x_k \in X_k$ represents a ball $B(x_k)$ of radius 2^k of U_1 , and by the properties of the ultrametric, this ball has diameter less or equal than 2^k . By the definition of γ_{ϱ_Φ} , if $\text{diam}(B(x_k)) \leq 2^k$ then $\text{diam}(\Phi(B(x_k))) \leq 2^{\gamma_{\varrho_\Phi}(k)} = 2^{\sigma(k)}$ and, since $Y_{\sigma(k)}$ is the partition of U_2 in balls of radius $2^{\sigma(k)}$, there is a unique point $y_{\sigma(k)} \in Y_{\sigma(k)}$ such that $\Phi(B(x_k)) \subset B(y_{\sigma(k)})$. Then, the map f_k such that $f_k(x_k) := y_{\sigma(k)}$ is well defined and it is surjective because Φ is surjective. It is immediate to check that the diagram commutes.

The morphism (f_k, σ) induces a map $\Phi: \text{end}(X_k, \Phi_k) \rightarrow \text{end}(Y_k, \Psi_k)$ where $\Phi((x_k))$ is the unique sequence $(y_k) \in \text{end}(Y_k, \Psi_k)$ such that $y_{\sigma(k)} = f_k(x_k)$. Now, for any $t \in (2^{k-1}, 2^k]$, $k \in \mathbb{Z}$, let $\varrho_\Phi(t) = 2^{\sigma(k)}$ and $\varrho_\Phi(0) = 0$. It is readily seen that ϱ_Φ is an extension function and Φ is an all scale uniform map. From Lemma 3.5 together with Proposition 3.2, it follows that there is an all scale uniform map $f: U_1 \rightarrow U_2$. $\square$

Lemma 3.7. *If $\Phi: X \rightarrow Y$ is an all scale uniform equivalence then there are increasing maps $\alpha(\Phi), \beta(\Phi^{-1}): \mathbb{Z} \rightarrow \mathbb{Z}$ such that $\gamma_{\varrho_\Phi}(\alpha(i)) \leq \beta(i)$, $\gamma_{\varrho_{\Phi^{-1}}}(\beta(i)) \leq \alpha(i+1)$ for every $i \in \mathbb{Z}$.*

Proof. First, let $\alpha(0) = 0$ and $\beta(0) = \gamma_\Phi(0)$.

If we have defined $\alpha(i-1), \beta(i-1)$ for any $i > 0$ then, it suffices make $\alpha(i) = \max\{\alpha(i-1) + 1, \gamma_{\Phi^{-1}}(\beta(i-1))\}$ and $\beta(i) = \max\{\beta(i-1) + 1, \gamma_\Phi(\alpha(i))\}$.

Since $\lim_{k \rightarrow -\infty} \gamma_{\Phi^{-1}}(k) = -\infty$ for any $\alpha(i+1)$ there exist some $k_{\alpha(i+1)}$ such that for any $k \leq k_{\alpha(i+1)}$, $\gamma_{\Phi^{-1}}(k) \leq \alpha(i+1)$.

Since $\lim_{k \rightarrow -\infty} \gamma_\Phi(k) = -\infty$ for any $\beta(i)$ there exist some $k_{\beta(i)}$ such that for any $k \leq k_{\beta(i)}$, $\gamma_\Phi(k) \leq \beta(i)$.

If we have defined $\alpha(i+1), \beta(i+1)$ for any $i < 0$ then, it suffices make $\beta(i) = \min\{\beta(i+1) - 1, k_{\alpha(i+1)}\}$ and $\alpha(i) = \min\{\alpha(i+1) - 1, k_{\beta(i)}\}$. $\square$

Proposition 3.8. *If $(X_k, \Phi_k), (Y_k, \Psi_k)$ are two D -chains, then $(X_k, \Phi_k) \sim_{z-z} (Y_k, \Psi_k)$ if and only if $\text{end}(X_k, \Phi_k)$ and $\text{end}(Y_k, \Psi_k)$ are all scale uniform equivalent.*

Proof. If (Z_k, ϕ_k) is a common zig-zag D -chain, it suffices to check that $\text{end}(X_k, \Phi_k)$ and $\text{end}(Y_k, \Psi_k)$ are all scale uniform equivalent to $\text{end}(Z_k, \phi_k)$ and this follows immediately from Lemma 3.5.

Now, suppose that there is an all scale uniform equivalence $\Phi: \text{end}(X_k, \Phi_k) \rightarrow \text{end}(Y_k, \Psi_k)$. By Lemma 3.7, there are increasing maps $\alpha(\Phi), \beta(\Phi^{-1}): \mathbb{Z} \rightarrow \mathbb{Z}$ such that $\gamma_{\varrho_\Phi}(\alpha(i)) \leq \beta(i)$, $\gamma_{\varrho_{\Phi^{-1}}}(\beta(i)) \leq \alpha(i+1)$ for every $i \in \mathbb{Z}$.

Therefore, Φ and Φ^{-1} canonically induce unique surjective maps $\mathcal{V}_{2i-1}: X_{\alpha(i)} \rightarrow Y_{\beta(i)}$ and $\mathcal{V}_{2i}: Y_{\beta(i)} \rightarrow X_{\alpha(i+1)}$. Since Φ is a bijection, $\mathcal{V}_{k+1} \circ \mathcal{V}_k$ coincides with the map induced by inclusion and therefore, making $Z_{2i-1} = X_{\alpha(i)}$ and $Z_{2i} = Y_{\beta(i)}$, $(Z_k, \mathcal{V}_k)$ is a common zig-zag D -chain of $(X_k, \Phi_k), (Y_k, \Psi_k)$. $\square$

From this, and Proposition 3.2, it follows:

Theorem 3.9. *Two complete ultrametric spaces are all scale uniform equivalent if and only if there is a common zig-zag D -chain between their associated D -chains.* ■

Let $(X_k, \Phi_k), (Y_k, \Psi_k)$ be two D -chains and $(Z_k, \mathcal{V}_k)$ a common zig-zag D -chain with increasing maps $\alpha, \beta: \mathbb{Z} \rightarrow \mathbb{Z}$ such that $Z_{2k-1} = X_{\alpha(k)}$ and $Z_{2k} = Y_{\beta(k)}$. Let us define

$$f_Z: \text{end}(X_{\alpha(k)}, \tilde{\Phi}_k) \rightarrow \text{end}(Y_{\beta(k)}, \tilde{\Psi}_k)$$

such that for any end point $(x_{\alpha(k)}) \in \text{end}(X_{\alpha(k)}, \tilde{\Phi}_k)$, $f_Z((x_{\alpha(k)})) = (\mathcal{V}_{2k-1}(x_{\alpha(k)})) \in \text{end}(Y_{\beta(k)}, \tilde{\Psi}_k)$.

Let us recall that a function between metric spaces $f: X \rightarrow Y$ is called *bi-Lipschitz* if there is a constant $K > 0$ such that for any pair of points $x, x' \in X$, $\frac{1}{K} \cdot d_X(x, x') \leq d_Y(f(x), f(x')) \leq K \cdot d_X(x, x')$. If there is such a map, we say that X, Y are bi-Lipschitz equivalent.

Proposition 3.10. *Given two D -chains $(X_k, \Phi_k), (Y_k, \Psi_k)$, then their end spaces $\text{end}(X_k, \Phi_k)$ and $\text{end}(Y_k, \Psi_k)$ are all scale uniform equivalent if and only if there are increasing sequences $\alpha, \beta: \mathbb{Z} \rightarrow \mathbb{Z}$ such that $\text{end}(X_{\alpha(k)}, \tilde{\Phi}_k)$ and $\text{end}(Y_{\beta(k)}, \tilde{\Psi}_k)$ are bi-Lipschitz equivalent.*

Proof. If $\text{end}(X_{\alpha(k)}, \tilde{\Phi}_k)$ and $\text{end}(Y_{\beta(k)}, \tilde{\Psi}_k)$ are bi-Lipschitz equivalent, then they are, in particular, all scale equivalent and so they are $\text{end}(X_n, \Phi_n)$ and $\text{end}(Y_n, \Psi_n)$ by Lemma 3.5.

If $\text{end}(X_k, \Phi_k)$ and $\text{end}(Y_k, \Psi_k)$ are all scale equivalent then, by Proposition 3.8, there is a zig-zag common D -chain $(Z_k, \mathcal{V}_k)$ defined by sequences α, β . The map $f_Z: \text{end}(X_{\alpha(k)}, \tilde{\Phi}_k) \rightarrow \text{end}(Y_{\beta(k)}, \tilde{\Psi}_k)$ defined above holds that for any pair of end points $(x_{\alpha(k)}), (y_{\alpha(k)}) \in \text{end}(X_{\alpha(k)}, \tilde{\Phi}_k)$, $D_{T(\beta)}(f_Z((x_{\alpha(k)})), f_Z((y_{\alpha(k)}))) \leq D_{T(\alpha)}(x_{\alpha(k)}, y_{\alpha(k)}) \leq 2 \cdot D_{T(\beta)}(f_Z((x_{\alpha(k)})), f_Z((y_{\alpha(k)})))$. ■

Corollary 3.11. *Two ultrametric spaces U_1, U_2 are asymorphic if and only if there are increasing sequences $\alpha, \beta: \mathbb{N} \rightarrow \mathbb{N}$ such that, for $(X_k, \Phi_k), (Y_k, \Psi_k)$ their associated D -chains, $\text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$ and $\text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$ are bi-Lipschitz equivalent.* ■

This can be translated into relations between ultrametric spaces avoiding D -chains. Proposition 2.2 in [2] states

Proposition 3.12. *Let (X, d) be a metric space. The metric d is an ultrametric if and only if $f(d)$ is a metric for every nondecreasing function $f: \mathbb{R}_+ \rightarrow \mathbb{R}_+$.* ■

In particular, the new metric is also an ultrametric. Given an ultrametric space (U, d) and a non-decreasing map $f: \mathbb{R}_+ \rightarrow \mathbb{R}_+$, let us denote this new ultrametric as d_f , where $d_f(x, y) := f(d(x, y))$.

Let (U, d) be an ultrametric space and an increasing map $\gamma: \mathbb{Z} \rightarrow \mathbb{Z}$. Let us define $f_\gamma: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ a non-decreasing function such that for any $t \in (2^{\gamma(k-1)}, 2^{\gamma(k)}]$ $f_\gamma(t) = 2^k$ for every k . Let us denote simply by $(U, d(\gamma))$ the ultrametric space (U, d_{f_γ}) which depends only on the original ultrametric and γ .

Thus, from Corollary 3.11 we obtain that,

Corollary 3.13. *Two ultrametric spaces $(U_1, d_1), (U_2, d_2)$ are asymorphic if and only if there are increasing maps $\gamma_1, \gamma_2: \mathbb{Z} \rightarrow \mathbb{Z}$ such that $(U_1, d_1(\gamma_1))$ and $(U_2, d_2(\gamma_2))$ are bi-Lipschitz equivalent.* ■

4. COARSE EQUIVALENCES

In this section we treat the category $\mathcal{C}_2$. All we do, is to consider only the right side of the chain in the previous section and adapt the construction in some technical details.

Given a bornologous multimap Φ and its expansion function ϱ_Φ , let us define $\gamma_{\varrho_\Phi} : \mathbb{N} \rightarrow \mathbb{N}$ as follows,

$$(2) \quad \gamma_{\varrho_\Phi}(n) := [\log_2(\varrho_\Phi(2^n))] + 1.$$

Hence, for all points $x, x' \in X$, if $d_X(x, x') \leq 2^n$ then $d_Y(f(x), f(x')) \leq 2^{\gamma_{\varrho_\Phi}(n)}$ and γ_{ϱ_Φ} is non-decreasing. We may assume, with no loss of generality, that $\lim_{t \rightarrow \infty} \varrho_\Phi(t) = \infty$ and therefore $\lim_{n \rightarrow \infty} \gamma_{\varrho_\Phi}(n) = \infty$.

Remark 4.1. *If Φ is an asyrmorphism between unbounded metric spaces then necessarily $\lim_{t \rightarrow \infty} \varrho_\Phi(t) = \infty$ since Φ^{-1} is a bornologous surjective map.*

There is a correspondence between ultrametric spaces and D_+ -chains. For each $n \in \mathbb{N}$ let X_n be the partition of U in closed balls of radius 2^n . For each $x_n \in X_n$ let us denote by $B(x_n)$ the associated closed ball in U . Let $\Phi_n : X_n \rightarrow X_{n+1}$ the map canonically induced by the inclusion for any $n \in \mathbb{N}$. (X_n, Φ_n) will be called the D_+ -chain associated to U . Conversely, given a D_+ -chain we can obtain an ultrametric space as follows.

The end space is:

$$\text{end}(X_n, \Phi_n) := \{(x_n)_{n \in \mathbb{N}} \mid x_n \in X_n \text{ and } \Phi(x_n) = x_{n+1}\},$$

and the metric

$$D((x_n), (y_n)) = 2^{n_0} \text{ where } n_0 = \min\{n : x_n = y_n\}.$$

D is well defined since $\lim_{\rightarrow} (X_n, \Phi_n)$ is trivial and $(\text{end}(X_n, \Phi_n), D)$ is an ultrametric space.

Proposition 4.2. *If U is an ultrametric space and (X_n, Φ_n) is the D_+ -chain associated to U , then U and $\text{end}(X_n, \Phi_n)$ are asyrmorphic (i.e., coarse equivalent).*

Proof. Consider the multi-map $\Phi : U \Rightarrow \text{end}(X_n, \Phi_n)$ where $\Phi := \{(x, (x_n)_{n \in \mathbb{N}}) \mid x \in B(x_1)\}$. Thus, if $\text{diam}(A) \leq 2$ then there exists some x_1 such that $A \subset B(x_1)$ and $\text{diam}(\Phi(A)) = 0$. If $2^n < \text{diam}(A) \leq 2^{n+1}$, then for each $x_n \in X_n$, $A \not\subset B(x_n)$ and, by the properties of the ultrametric, there is a unique $x_{n+1} \in X_{n+1}$ such that $A \subset B(x_{n+1})$ and therefore, $\text{diam}(\Phi(A)) = 2^{n+1}$ for every $n > 1$. Hence, Φ is an asyrmorphism. $\square$

Definition 4.3. A morphism of D_+ -chains $(f_n, \sigma) : (X_n, \Phi_n) \rightarrow (Y_n, \Psi_n)$ consists of a non-decreasing function $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ such that $\lim_{n \rightarrow \infty} \sigma(n) = \infty$, and maps $f_n : X_n \rightarrow Y_{\sigma(n)}$ such that the following diagram commutes:

$$\begin{array}{ccccccc} X_1 & \longrightarrow & X_2 & \longrightarrow & X_3 & \longrightarrow & \dots \\ & \searrow f_1 & & \searrow f_2 & & \searrow f_3 & \\ & Y_{\sigma(1)} & \longrightarrow & \dots & \longrightarrow & Y_{\sigma(2)} & \longrightarrow \dots \end{array}$$

Lemma 4.4. *If (X_n, Φ_n) is a D_+ -chain, $\alpha : \mathbb{N} \rightarrow \mathbb{N}$ is an increasing map and $(X_{\alpha(n)}, \tilde{\Phi}_n)$ is the α -subchain, then $\text{end}(X_n, \Phi_n)$ is asyrmorphic to $\text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$.*

Proof. There is a canonical map $i: \text{end}(X_n, \Phi_n) \rightarrow \text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$ with $i((x_n)) = (x_{\alpha(n)})$.

Let us define the function $\Lambda: \mathbb{N} \rightarrow \mathbb{N}$ such that $\Lambda(n) = \min\{k : \alpha(k) \geq n\}$. Since α is increasing, it follows that Λ is non-decreasing and $\lim_{n \rightarrow \infty} \Lambda(n) = \infty$. Now, consider the function $\Gamma: [0, \infty) \rightarrow [0, \infty)$ such that $\Gamma(x) = 2^{\Lambda(1)}$ for $x \leq 1$ and for any $x \in (2^{n-1}, 2^n]$, $\Gamma(x) = 2^{\Lambda(n)}$, $n \in \mathbb{N}$. Clearly Γ is non-decreasing and $\lim_{x \rightarrow \infty} \Gamma(x) = \infty$. For any pair of end points $(x_n), (y_n) \in \text{end}(X_n, \Phi_n)$ with $D((x_n), (y_n)) = 2^{n_0}$ the distance between their correspondent subsequences $i(x_n) = (x_{\alpha(n)}), i(y_n) = (y_{\alpha(n)})$ is exactly $D'((x_{\alpha(n)}), (y_{\alpha(n)})) = \Gamma(2^{n_0}) = 2^{\Lambda(n_0)}$. Then Γ is an extension function for $i: \text{end}(X_n, \Phi_n) \rightarrow \text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$ and i is bornologous. A similar argument works for the multi-map i^{-1} . $\square$

Proposition 4.5. *Consider two ultrametric spaces U_1, U_2 and $(X_n, \Phi_n), (Y_n, \Psi_n)$ their associated D_+ -chains. Then, there is a bornologous multi-map $\Phi: U_1 \Rightarrow U_2$ if and only if there is a morphism of D_+ -chains $(f_n, \sigma): (X_n, \Phi_n) \rightarrow (Y_n, \Psi_n)$.*

Proof. If there is a bornologous multi-map $\Phi: U_1 \Rightarrow U_2$, consider the map $\gamma_{\Phi}: \mathbb{N} \rightarrow \mathbb{N}$ from (2). Making $\sigma = \gamma_{\Phi}$, Φ canonically induces the maps $f_n: X_n \rightarrow Y_{\sigma(n)}$ and the diagram commutes.

The morphism (f_n, α) induces a multi-map $\Phi: \text{end}(X_n, \Phi_n) \Rightarrow \text{end}(Y_n, \Psi_n)$ where $\Phi((x_n))$ is the set of sequences $(y_n) \in \text{end}(Y_n, \Psi_n)$ such that $y_{\sigma(n)} = f_n(x_n)$. It is immediate to check that this multi-map is bornologous, and from Lemma 4.4 together with Proposition 4.2, it follows that there is a bornologous multi-map $\Phi: U_1 \Rightarrow U_2$. $\square$

Lemma 4.6. *If $\Phi: X \Rightarrow Y$ is an asymorphism then there are increasing maps $\alpha(\Phi), \beta(\Phi^{-1}): \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ such that $\gamma_{\Phi}(\alpha(i)) \leq \beta(i)$, $\gamma_{\Phi^{-1}}(\beta(i)) \leq \alpha(i+1)$ for every $i \in \mathbb{Z}_+$.*

Proof. First, let $\alpha(1) = 1$ and $\beta(1) = \gamma_{\Phi}(1)$.

If we have defined $\alpha(i-1), \beta(i-1)$ for any $i > 1$ then, it suffices make $\alpha(i) = \max\{\alpha(i-1) + 1, \gamma_{\Phi^{-1}}(\beta(i-1))\}$ and $\beta(i) = \max\{\beta(i-1) + 1, \gamma_{\Phi}(\alpha(i))\}$. $\square$

Proposition 4.7. *If $(X_n, \Phi_n), (Y_n, \Psi_n)$ are two D_+ -chains, then $(X_n, \Phi_n) \sim_{z-z} (Y_n, \Psi_n)$ if and only if $\text{end}(T, v)$ and $\text{end}(T', w)$ are asymorphic.*

Proof. If there is a common zig-zag chain, the existence of an asymorphism follows immediately from Lemma 4.4.

Now, suppose that there is an asymorphism $\Phi: \text{end}(X_n, \Phi_n) \Rightarrow \text{end}(Y_n, \Psi_n)$. By Lemma 4.6, if Φ is an asymorphism then there are increasing maps $\alpha(\Phi), \beta(\Phi^{-1}): \mathbb{N} \rightarrow \mathbb{N}$ such that $\gamma_{\Phi}(\alpha(i)) \leq \beta(i)$, $\gamma_{\Phi^{-1}}(\beta(i)) \leq \alpha_{i+1}$ for every $i \in \mathbb{N}$.

Therefore, Φ and Φ^{-1} canonically induce (as we saw in 3.8) unique surjective maps $\mathcal{V}_{2i-1}: X_{\alpha(i)} \rightarrow Y_{\beta(i)}$ and $\mathcal{V}_{2i}: Y_{\beta(i)} \rightarrow X_{\alpha(i+1)}$. The ball associated to the vertex $\mathcal{V}_{k+1} \circ \mathcal{V}_k(x)$ will contain, by construction, the ball associated to the vertex x , and $\mathcal{V}_{k+1} \circ \mathcal{V}_k$ coincides with the map induced by the inclusion. Therefore, making $Z_{2i-1} = X_{\alpha(i)}$ and $Z_{2i} = Y_{\beta(i)}$, $(Z_n, \mathcal{V}_n)$ is a common zig-zag D_+ -chain of $(X_n, \Phi_n), (Y_n, \Psi_n)$. $\square$

From this, and Proposition 4.2, it follows:

Theorem 4.8. *Two ultrametric spaces are coarse equivalent if and only if there is a common zig-zag chain between their associated D_+ -chains.* $\blacksquare$

Let $(X_n, \Phi_n), (Y_n, \Psi_n)$ be two D_+ -chains and $(Z_n, \mathcal{V}_n)$ a common zig-zag D_+ -chain with increasing maps $\alpha, \beta: \mathbb{N} \rightarrow \mathbb{N}$ such that $Z_{2n-1} = X_{\alpha(n)}$ and $Z_{2n} = Y_{\beta(n)}$. Let us define

$$f_Z: \text{end}(X_{\alpha(n)}, \tilde{\Phi}_n) \rightarrow \text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$$

such that for any end point $(x_{\alpha(n)}) \in \text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$, $f_Z((x_{\alpha(n)})) = (\mathcal{V}_{2n-1}(x_{\alpha(n)})) \in \text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$.

Proposition 3.10 is not true in the case of D_+ -chains, since the induced map between the end spaces is not necessarily injective. There would be a bi-Lipschitz equivalence restricted to large scale. Moreover,

Proposition 4.9. *Given two D_+ -chains $(X_n, \Phi_n), (Y_n, \Psi_n)$, then $\text{end}(X_n, \Phi_n)$ and $\text{end}(Y_n, \Psi_n)$ are asyomorphic if and only if there are increasing sequences $\alpha, \beta: \mathbb{N} \rightarrow \mathbb{N}$ and a map $F: \text{end}(X_{\alpha(n)}, \tilde{\Phi}_n) \rightarrow \text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$ such that for any pair of end points $(x_{\alpha(n)}), (y_{\alpha(n)}) \in \text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$,*

$$D_{T(\beta)}(f((x_{\alpha(n)})), f((y_{\alpha(n)}))) \leq D_{T(\alpha)}((x_{\alpha(n)}), (y_{\alpha(n)}))$$

and if $D_{T(\alpha)}((x_{\alpha(n)}), (y_{\alpha(n)})) > 2$, then

$$D_{T(\alpha)}((x_{\alpha(n)}), (y_{\alpha(n)})) \leq 2 \cdot D_{T(\beta)}(f((x_{\alpha(n)})), f((y_{\alpha(n)}))).$$

Proof. If there is such a map F , then $\text{end}(X_{\alpha(n)}, \tilde{\Phi}_n), \text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$ are, in particular, asyomorphic and so they are $\text{end}(X_n, \Phi_n)$ and $\text{end}(Y_n, \Psi_n)$ by Lemma 4.4.

If $\text{end}(X_n, \Phi_n)$ and $\text{end}(Y_n, \Psi_n)$ are asyomorphic then, by Proposition 4.7, there is a common zig-zag D_+ -chain given by sequences α, β . It is immediate to check that the map $f_Z: \text{end}(X_{\alpha(n)}, \tilde{\Phi}_n) \rightarrow \text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$ holds the conditions above. $\square$

5. UNIFORM HOMEOMORPHISMS

In this section we treat the category $\mathcal{C}_3$. The idea is to consider only the left side of the D -chain but to avoid using as index set the negative integers we change the orientation of the chain and therefore the construction of γ . Also, we have to be careful with the fact that the expansion function is defined on some interval $[0, S]$. Let $\mathbb{N}_{\geq n_0} := \{n \in \mathbb{N} \mid n \geq n_0\}$.

Given a uniformly continuous map Φ and its expansion function $\varrho_\Phi: [0, S] \rightarrow [0, \frac{1}{2}]$ let $n_0 \in \mathbb{N}$ such that $2^{-n_0} \leq S < 2^{-n_0+1}$ if $S < \frac{1}{2}$ or $n_0 = 1$ otherwise, and let us define $\gamma_{\varrho_\Phi}: \mathbb{N}_{\geq n_0} \rightarrow \mathbb{N}$ as follows,

$$(3) \quad \gamma_{\varrho_\Phi}(n) := \lceil -\log_2(\varrho_\Phi(2^{-n})) \rceil.$$

Hence, for all points $x, x' \in X$, if $d_X(x, x') \leq 2^{-n}$ then $d_Y(\Phi(x), \Phi(x')) \leq 2^{-\gamma_{\varrho_\Phi}(n)}$ and γ_{ϱ_Φ} is non-decreasing, $\lim_{n \rightarrow \infty} \gamma_{\varrho_\Phi}(n) = \infty$ since Φ is uniformly continuous.

Given an ultrametric space U there is a D_- -chain associated to it. For each $n \in \mathbb{N}$ let X_n be the partition of U in balls of radius 2^{-n} . Let $\Phi_n: X_{n+1} \rightarrow X_n$ the map canonically induced by the inclusion for any $n \in \mathbb{N}$. (X_n, Φ_n) will be called the D_- -chain associated to U . Conversely, given a D_- -chain we can obtain an ultrametric space as follows.

The end space is then:

$$\text{end}(X_n, \Phi_n) := \{(x_n)_{n \in \mathbb{N}} \mid x_n \in X_n \text{ and } \Phi(x_{n+1}) = x_n\},$$

and let the metric be

$$D((x_n), (y_n)) = \begin{cases} 2^{-n_0} & \text{if there is } n_0 = \max\{n : x_n = y_n\}, \\ 1 & \text{if } x_n \neq y_n \ \forall n. \end{cases}$$

Clearly, D is an ultrametric.

Proposition 5.1. *If (U, d) is a complete ultrametric space and (X_n, Φ_n) is the D_- -chain associated to U , then (U, d) and $\text{end}(X_n, \Phi_n)$ are uniformly homeomorphic.*

Proof. First, note that if (U, d) is complete there is a bijection $i: U \rightarrow \text{end}(X_n, \Phi_n)$.

Notice that, by the properties of the ultrametric, $d(x, y) \leq 2^{-n}$ if and only if the points are in the same ball in the partition X_n . Hence, if $d(x, y) \leq 2^{-n}$, $d(x, y) \leq D(i(x), i(y)) \leq 2d(x, y)$ and i is a uniform homeomorphism. $\square$

In the case of uniform maps, we need to consider in the description of the morphisms of D_- chains the radius such that the image of the ball will have diameter bounded by $1/2$ (i.e. the interval $[0, S]$ on which the expansion function is defined):

Definition 5.2. A morphism of D_- -chains $(f_n, \sigma, n_0): (X_n, \Phi_n) \rightarrow (Y_n, \Psi_n)$ consists of a natural number n_0 , a non-decreasing function $\sigma: \mathbb{N}_{\geq n_0} \rightarrow \mathbb{N}$ such that $\lim_{n \rightarrow \infty} \sigma(n) = \infty$, and maps $f_n: X_n \rightarrow Y_{\sigma(n)} \ \forall n \geq n_0$ such that the following diagram commutes:

$$\begin{array}{ccccccc} \longleftarrow & & X_n & & \longleftarrow & & X_{n+1} & & \longleftarrow & & X_{n+2} \\ & \searrow f_n & & \searrow f_{n+1} & & \searrow f_{n+2} & & & & & \\ Y_{\sigma(n)} & \longleftarrow & \dots & \longleftarrow & Y_{\sigma(n+1)} & \longleftarrow & \dots & \longleftarrow & Y_{\sigma(n+2)} & \longleftarrow & \end{array}$$

Lemma 5.3. *If (X_n, Φ_n) is a D_- -chain, $\alpha: \mathbb{N} \rightarrow \mathbb{N}$ is an increasing map and $(X_{\alpha(n)}, \tilde{\Phi}_n)$ is the α -subchain, then $\text{end}(X_n, \Phi_n)$ is uniformly homeomorphic to $\text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$.*

Proof. Consider the canonical map $i: \text{end}(X_n, \Phi_n) \rightarrow \text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$.

Let us define the function $\Lambda: \mathbb{N} \rightarrow \mathbb{N}$ such that $\Lambda(n) = \max\{k : \alpha(k) \leq n\}$. Since α is increasing, it follows that Λ is non-decreasing and $\lim_{n \rightarrow \infty} \Lambda(n) = \infty$. Now, consider the function $\Gamma: [0, \infty) \rightarrow [0, \infty)$ such that $\Gamma(S) = \frac{1}{2}$ and for any $x \in [S \cdot 2^{-n}, S \cdot 2^{-n+1})$, $\Gamma(x) = 2^{-\Lambda(n)}$. Clearly Γ is non-decreasing, $\lim_{x \rightarrow 0} \Gamma(x) = 0$. For any two end points $(x_n), (y_n) \in \text{end}(X_n, \Phi_n)$ with $D((x_n), (y_n)) = 2^{-n_0}$ the distance between their correspondent subsequences $i(x_n) = (x_{\alpha(n)}), i(y_n) = (y_{\alpha(n)})$ is exactly $D'((x_{\alpha(n)}), (y_{\alpha(n)})) = \Gamma(2^{-n_0}) = 2^{-\Lambda(n_0)}$. Then Γ is an extension function for $i: \text{end}(X_n, \Phi_n) \rightarrow \text{end}((X_{\alpha(n)}, \tilde{\Phi}_n))$ and i is a uniform homeomorphism. A similar argument works for i^{-1} . $\square$

Proposition 5.4. *Consider two complete ultrametric spaces U_1, U_2 and let $(X_n, \Phi_n), (Y_n, \Psi_n)$ be their associated D_- -chains. Then, there is a uniformly continuous map $\Phi: U_1 \rightarrow U_2$ if and only if there is a morphism of D_- -chains $(f_n, \alpha, n_0): (X_n, \Phi_n) \rightarrow (Y_n, \Psi_n)$.*

Proof. If there is a uniformly continuous map $\Phi: U_1 \rightarrow U_2$, consider the map $\gamma_{\Phi}: \mathbb{N} \rightarrow \mathbb{N}$ from (3). Making $\sigma = \gamma_{\Phi}$, Φ canonically induces the maps $f_n: X_n \rightarrow Y_{\sigma(n)} \ \forall n \geq n_0$ and the diagram commutes.

The morphism (f_n, α, n_0) induces a map $F: \text{end}(X_n, \Phi_n) \rightarrow \text{end}(Y_n, \Psi_n)$ where $F((x_n))$ is the unique sequence $(y_n) \in \text{end}(Y_n, \Psi_n)$ such that $y_{\sigma(n)} = f_n(x_n) \ \forall n \geq$

n_0 . It is immediate to check that this map is uniformly continuous, and from Lemma 5.3 together with Proposition 5.1, it follows that there is a uniformly continuous map $\Phi : U_1 \rightarrow U_2$. $\square$

Lemma 5.5. *If $\Phi : X \rightarrow Y$ is a uniform homeomorphism then there are increasing maps $\alpha(\Phi), \beta(\Phi^{-1}) : \mathbb{N} \rightarrow \mathbb{N}$ such that $\gamma_{\varrho_{\Phi^{-1}}}(\beta(i)) \leq \alpha(i)$, $\gamma_{\varrho_{\Phi}}(\alpha(i+1)) \leq \beta(i)$ for every $i \in \mathbb{N}$.*

Proof. First, let $\alpha(1) = 1$.

Since $\lim_{n \rightarrow \infty} \gamma_{\Phi^{-1}}(n) = \infty$ for any $\alpha(i)$ there exist some $n_{\alpha(i)} > 0$ such that for any $n \geq n_{\alpha(i+1)}$, $\gamma_{\Phi^{-1}}(n) \geq \alpha(i)$.

Let $\beta(1) = n_{\alpha(1)}$.

Since $\lim_{n \rightarrow \infty} \gamma_{\Phi}(n) = \infty$ for any $\beta(i)$ there exist some $n_{\beta(i)}$ such that for any $n \geq n_{\beta(i)}$, $\gamma_{\Phi}(n) \geq \beta(i)$.

If we have defined $\alpha(i-1), \beta(i-1)$, then, it suffices make $\alpha(i) = \max\{\alpha(i-1) + 1, n_{\beta(i-1)}\}$ and $\beta(i) = \max\{\beta(i-1) + 1, n_{\alpha(i)}\}$. $\square$

Proposition 5.6. *If $(X_n, \Phi_n), (Y_n, \Psi_n)$ are two D_- -chains, then $(X_n, \Phi_n) \sim_{z-z} (Y_n, \Psi_n)$ if and only if $\text{end}(T, v), \text{end}(T', w)$ are uniform homeomorphic.*

Proof. If there is a common zig-zag D_- -chain, the existence of an asyomorphism follows immediately from Lemma 5.3.

Now, suppose there is a uniform homeomorphism $\Phi : \text{end}(X_n, \Phi_n) \rightarrow \text{end}(Y_n, \Psi_n)$. Then, by Lemma 5.5, there are increasing maps $\alpha(\Phi), \beta(\Phi^{-1}) : \mathbb{N} \rightarrow \mathbb{N}$ such that $\gamma_{\varrho_{\Phi^{-1}}}(\beta(i)) \leq \alpha(i)$, $\gamma_{\varrho_{\Phi}}(\alpha(i+1)) \leq \beta(i)$ for every $i \in \mathbb{N}$.

Therefore, Φ and Φ^{-1} induce respectively unique surjective maps $\mathcal{V}_{2i-1} : Y_{\beta(i)} \rightarrow X_{\alpha(i)}$ and $\mathcal{V}_{2i} : X_{\alpha(i+1)} \rightarrow Y_{\beta(i)}$. $\mathcal{V}_{k+1} \circ \mathcal{V}_k$ coincides with the map induced by the inclusion, and hence, making $Z_{2i-1} = X_{\alpha(i)}$ and $Z_{2i} = Y_{\beta(i)}$, $(Z_n, \mathcal{V}_n)$ is a common zig-zag D_- -chain of $(X_n, \Phi_n), (Y_n, \Psi_n)$. $\square$

From this, and Proposition 5.1, it follows:

Theorem 5.7. *Two complete ultrametric spaces U_1, U_2 are uniformly homeomorphic if and only if there is a common zig-zag D_- -chain between their associated D_- -chains.* $\blacksquare$

Definition 5.8. *A function between metric spaces $f : X \rightarrow Y$ is small scale bi-Lipschitz if there is a constant $K > 0$ and a real number $\epsilon > 0$ such that for any pair of points $x, x' \in X$ with $d(x, x') < \epsilon$, $\frac{1}{K} \cdot d_X(x, x') \leq d_Y(f(x), f(x')) \leq K \cdot d_X(x, x')$. In there is such a map, we say that X, Y are small scale bi-Lipschitz equivalent.*

Let $(X_n, \Phi_n), (Y_n, \Psi_n)$ be two D_- -chains and $(Z_n, \mathcal{V}_n)$ a common zig-zag D_- -chain with increasing maps $\alpha, \beta : \mathbb{N} \rightarrow \mathbb{N}$ such that $Z_{2n-1} = X_{\alpha(n)}$ and $Z_{2n} = Y_{\beta(n)}$. Let us define

$$f_Z : \text{end}(X_{\alpha(n)}, \tilde{\Phi}_n) \rightarrow \text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$$

such that for any end point $(x_{\alpha(n)}) \in \text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$, $f_Z((x_{\alpha(n)}))$ is the unique end point $(y_{\beta(n)}) \in \text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$ such that $\mathcal{V}_{2n-2}(x_{\alpha(n)}) = y_{\beta(k-1)} \forall n \geq 2$.

Proposition 5.9. *Given two D_- -chains $(X_n, \Phi_n), (Y_n, \Psi_n)$, then $\text{end}(X_n, \Phi_n)$ and $\text{end}(Y_n, \Psi_n)$ are uniformly homeomorphic if and only if there are increasing sequences $\alpha, \beta : \mathbb{N} \rightarrow \mathbb{N}$ such that $\text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$ and $\text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$ are small scale bi-Lipschitz equivalent.*

Proof. If $\text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$ and $\text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$ are small scale bi-Lipschitz equivalent, then they are, in particular uniformly homeomorphic and so they are $\text{end}(X_n, \Phi_n)$ and $\text{end}(Y_n, \Psi_n)$ by Lemma 5.3.

If $\text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$ and $\text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$ are uniformly homeomorphic then, by Proposition 5.6, there is a common zig-zag D_- -chain given by sequences α, β . Then, the map $f_Z: \text{end}(X_{\alpha(n)}, \tilde{\Phi}_n) \rightarrow \text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$ described above, for any pair of end points $(x_{\alpha(n)}, (x'_{\alpha(n)})) \in \text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$ with $D_{T(\alpha)}((x_{\alpha(n)}, (x'_{\alpha(n)}))) \leq \frac{1}{4}$, holds that

$$D_{T(\alpha)}((x_{\alpha(n)}, (x'_{\alpha(n)}))) \leq D_{T(\beta)}(f_Z((x_{\alpha(n)}, (x'_{\alpha(n)})))) \leq 2 \cdot D_{T(\alpha)}((x_{\alpha(n)}, (x'_{\alpha(n)})))$$

and it is small scale bi-Lipschitz. $\square$

Corollary 5.10. *Two ultrametric spaces U_1, U_2 are uniformly homomorphic if and only if there are increasing sequences $\alpha, \beta: \mathbb{N} \rightarrow \mathbb{N}$ such that $\text{end}(X_{\alpha(n)}, \tilde{\Phi}_n)$ and $\text{end}(Y_{\beta(n)}, \tilde{\Psi}_n)$ are small scale bi-Lipschitz equivalent.* $\blacksquare$

Let (U, d) be an ultrametric space and $(n_i), i > 0$ an increasing sequence of numbers. Let us define $f_{(n_i)}: [0, \infty) \rightarrow \mathbb{N}$ a non-decreasing function such that for any $t \geq 2^{-n_1}$, $f(t) = 2^{-1}$ and for any $t \in (2^{-n_{i+1}}, 2^{-n_i}]$ $f(t) = 2^{-i}$ for every $i > 1$. Let us denote simply by $(U, d(n_i))$ the ultrametric space $(U, d_{f_{(n_i)}})$ which depends only on the original ultrametric and the sequence (n_i) .

Corollary 5.11. *Two ultrametric spaces $(U_1, d_1), (U_2, d_2)$ are uniformly homeomorphic if and only if there are increasing sequences of numbers $(n_i), (m_i)$ such that $(U_1, d_1(n_i))$ and $(U_2, d_2(m_i))$ are small scale bi-Lipschitz equivalent.* $\blacksquare$

6. TOWERS AND ADMISSIBLE MORPHISMS

In [1], Taras Banach and Ihor Zarichnyy give a classification of ultrametric spaces up to coarse geometry. They prove their results by induction on partially ordered sets called towers. The following definitions are stated as they appear in their paper.

A partially ordered set T is a *tree* if T has the smallest element and for every point $x \in T$ the lower cone $\downarrow x$ is well-ordered. By the *lower cone* (resp. *upper cone*) of a point x of a partially ordered set T we understand the set $\downarrow x = \{y \in T : y \leq x\}$ (resp. $\uparrow x = \{y \in T : y \geq x\}$). A subset A will be called a *lower* (resp. *upper*) *set* if $\downarrow a \subset A$ (resp. $\uparrow a \subset A$) for every $a \in A$. A partially ordered set T is *well-founded* if each subset $A \subset T$ has a minimal element $a \in A$. The minimality of a means that each point $a' \in A$ with $a' \leq a$ is equal to a . By $\min T$ we shall denote the set of all minimal elements of T .

Definition 6.1. *A partially ordered set T is called a tower if*

- (1) *T is well-founded;*
- (2) *any two elements $x, y \in T$ have the smallest upper bound $\sup\{x, y\}$ in T ;*
- (3) *for any $x \in T$ the upper cone $\uparrow x$ is linearly ordered;*
- (4) *for any point $a \in T$ there is a finite number $n = \text{lev}_T(a)$ such that for every minimal element $x \in \downarrow a$ of T the order interval $[x, a] = \uparrow x \cap \downarrow a$ has cardinality $|[x, a]| = n$.*

The function $\text{lev}_T: T \rightarrow \mathbb{N}$, $\text{lev}_T: a \mapsto \text{lev}_T(a)$, from the last item is called the *level function*.

The level function $lev_T: T \rightarrow \mathbb{N}$ divides T into the levels $L_i = lev_T^{-1}(i)$, $i \in \mathbb{N}$. The level $L_1 = \min T$ is called *the base* of T and denoted by $[T]$.

Each tower carries a canonic *path metric* d_T defined by the formula

$$d_T(x, y) = 2 \cdot lev_T(\sup(x, y)) - (lev_T(x) + lev_T(y)) \text{ for } x, y \in T.$$

The path metric restricted to the base $[T]$ of T is an ultrametric.

Given a tower T with levels L_i , we can define a D_+ -chain (L_i, Φ_i) with $\Phi: L_i \rightarrow L_{i+1}$ such that $\Phi(x_i) = x_{i+1}$ for any $x_i \leq x_{i+1}$.

Proposition 6.2. *For any tower T with levels L_i , $\text{end}(L_i, \Phi_i)$ is coarse equivalent to $[T]$.*

Proof. This is readily seen since $D(x, y) = 2^n$ if and only if $d_T(x, y) = 2n$. $\square$

For every point $x \in T$ of a tower T , the set $L_i \cap \downarrow x$ with $i = lev_T(x) - 1$ is denoted $\text{pred}(x)$ and it is called the set of *parents* of x .

Definition 6.3. *Let T_1, T_2 be two towers. A map $\phi: A \rightarrow T_2$ defined on a lower subset $A = \downarrow A$ of T_1 is called an admissible morphism if*

- (1) $lev(\phi(a)) = lev(a)$ for all $a \in A$;
- (2) $a \leq a'$ in A implies $\phi(a) \leq \phi(a')$;
- (3) $\phi(a) = \phi(a')$ for $a, a' \in A$ implies that $a, a' \in \text{pred}(v)$ for some $v \in T_1$;
- (4) $\phi(A)$ is a lower subset of T_2 ;
- (5) $|\phi(\max A)| \leq 1$, where $\max A$ stands for the (possibly empty) set of maximal elements of the domain A .

As we mentioned in the introduction, see 1.14, Lemma 2 in [1] states that the restriction to the base of an admissible morphism between towers is an asymorphism. Using D_+ -chains we proof that this is in fact an if and only if condition.

Consider two towers T, T' and their corresponding D_+ -chains $(X_n, \Phi_n), (Y_n, \Psi_n)$. Let $(Z_n, \mathcal{V}_n)$ be a common zig-zag D_+ -chain for $(X_n, \Phi_n), (Y_n, \Psi_n)$ with increasing maps $\alpha, \beta: \mathbb{N} \rightarrow \mathbb{N}$ such that $Z_{2n-1} = X_{\alpha(n)}$ and $Z_{2n} = X_{\beta(n)}$. $(X_{\alpha(n)}, \tilde{\Phi}_n), (Y_{\beta(n)}, \tilde{\Psi}_n)$ define subchains. Let $T(\alpha), T'(\beta)$ be the corresponding subtowers of T, T' defined respectively by levels $\alpha(n)$ and $\beta(n)$, $n \in \mathbb{N}$ and let $T(\alpha(n))$ denote $lev_{T(\alpha)}^{-1}(n)$ (its n^{th} level).

Let $f_Z: T(\alpha) \rightarrow T'(\beta)$ be such that for every n , $f_Z|_{T(\alpha(n))} = \mathcal{V}_{2n-1}: T(\alpha(n)) \rightarrow T'(\beta(n))$.

It is immediate to check the following:

Proposition 6.4. *Given a common zig-zag D_+ -chain (Z_n, r_n) for two towers T, T' , f_Z is an admissible map.* $\blacksquare$

From Proposition 6.4 together with 4.7 and 6.2, and Proposition 1.14 we conclude that

Corollary 6.5. *Given two towers T_1, T_2 , $[T_1]$ and $[T_2]$ are asymorphic if and only if there is an admissible map $f: T(\alpha) \rightarrow T'(\beta)$ for some pair of sequences α, β .* $\blacksquare$

What follows is a version of 4.9 for the metric given here to the base.

Proposition 6.6. *Given two towers T_1, T_2 , $[T_1]$ and $[T_2]$ are asymorphic if and only if there are increasing sequences $\alpha, \beta: \mathbb{N} \rightarrow \mathbb{N}$ such that $[T_1(\alpha)]$ and $[T_2(\beta)]$ are roughly isometric.*

Proof. If $[T_1(\alpha)]$ and $[T_2(\beta)]$ are roughly isometric, then they are, in particular asyomorphic. Proposition 6.2 and Lemma 4.4 yield that $[T_1]$ and $[T_2]$ are asyomorphic.

From Corollary 6.5, we obtain an admissible map $f: T(\alpha) \rightarrow T'(\beta)$. For any pair of points $x, y \in [T_1(\alpha)]$, condition (3) in the definition of admissible map implies that $d_{T(\beta)}(f(x), f(y)) \leq d_{T(\alpha)}(x, y) \leq d_{T(\beta)}(f(x), f(y)) + 2$. $\square$

REFERENCES

- [1] Banach, T., Zarichnyy, I. *The coarse classification of homogeneous ultra-metric spaces*. Preprint (arXiv:0801.2132).
- [2] Brodskiy, N. et al. *Dimension zero at all scales*. Topology and its Applications. 154 (2007) 2729–2740.
- [3] Buyalo, S., Schroeder, V. *Elements of Asymptotic Geometry*. EMS Monographs in Mathematics. Germany (2007).
- [4] Hughes, B. *Trees and ultrametric spaces: a categorial equivalence*. Advances in Mathematics. **189**,(2004)148-191.
- [5] Kühn K. *Direct Limits of Diagonal Chains of Type O, U, and Sp, and Their Homotopy Groups*. Communications in Algebra. 74 75–87 (2006).
- [6] Mardešić, S., Segal, J. *Shape theory*. North-Holland (1982).
- [7] Martínez-Perez, A. and Morón, M.A. *Uniformly continuous maps between ends of $\mathbb{R}$ -trees*. Math. Z. (To appear).
- [8] Martínez-Perez, A. and Morón, M.A. *Inverse sequences, rooted trees and their end spaces*. arXiv:0710.3004 [math.GT]
- [9] Roe, J. *Lectures on coarse geometry*. University Lecture Series, **vol.31** American Mathematical Society (2003).

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, UNIVERSIDAD COMPLUTENSE DE MADRID,
MADRID 28040 SPAIN

E-mail address: alvaro_martinez@mat.ucm.es